

Medicare Preventive Screening

Human Immunodeficiency Virus (HIV) Screening

(NCD 210.7)

What's Changed? No 2020 fourth quarter changes

HCPCS/CPT Codes

- 80081 —
Obstetric panel (includes HIV testing)
- G0432 —
Infectious agent antibody detection by enzyme immunoassay (eia) technique, hiv-1 and/or hiv-2, screening
- G0433 —
Infectious agent antibody detection by enzyme-linked immunosorbent assay (ELISA) technique, hiv-1 and/or hiv-2, screening
- G0435 —
Infectious agent antibody detection by rapid antibody test, hiv-1 and/or hiv-2, screening
- G0475 —
HIV antigen/antibody, combination assay, screening

ICD-10 Codes

Increased risk factors not reported: Z11.4

Increased risk factors reported: Z11.4 and Z72.51, Z72.52, Z72.53, or Z72.89

Pregnant patients: Z11.4 and Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, O09.90, O09.91, O09.92, or O09.93

NOTE: Additional ICD-10 codes may apply. See the [CMS ICD-10 webpage](#) for individual Change Requests (CRs) and the specific ICD-10-CM codes Medicare covers for this service, and [Contact your MAC](#) for guidance.

Medicare Covers

Certain patients without regard to perceived risk or at increased risk for HIV infection, including anyone who asks for the test, or pregnant women.

NOTE: The [Medicare National Coverage Determinations Manual, Chapter 1, Part 4, Section 210.7](#) defines “Increased risk for HIV infection.”

Medicare Preventive Screening

Frequency

- Annually for patients between the ages of 15 and 65 without regard to perceived risk
- Annually for patients younger than 15 and adults older than 65 at increased risk for HIV infection
- For pregnant patients, 3 times per pregnancy
 - First, when diagnosed as pregnant
 - Second, during the third trimester
 - Third, at labor, if ordered by their clinician

Patient Pays

- Copayment/Coinsurance waived
- Deductible waived

Other Notes

- For more information, refer to [Screening for the Human Immunodeficiency Virus \(HIV\) Infection](#).